performance evaluation four

ADULT FIRST AID - SEVERE, LIFE-THREATENING EXTERNAL BLEEDING



Student Name	
	Class D-t-
	Class Date

You are a trained first aid provider responding to a call received over your workplace two-way radio for a "person hit by gunfire." The shooter has fled. The scene is safe. You see a person sitting on the floor holding their hand over a bloody wound on their thigh. You have appropriate PPE and a first aid kit with a 10-pack of sterile 4" x 4" gauze sponges, but no hemostatic dressings or a tourniquet. Demonstrate what actions you would take next.

Procedure	Provider Action (Performance Criteria)	Instructor Prompt	Chec
Performs Assessment	 Assesses scene safety. Takes standard precautions (at a minimum, puts on disposable gloves). Assesses responsiveness. Activates EMS and/or EAP. After activating, and unless they are readily available, sends someone to get the first aid kit and an AED. 	"Shooter has fled. The scene is safe." "Person is responsive." "EMS/EAP activated." "Other first aid providers are on the way, bringing an AED and additional first aid kits."	
Assesses Breathing	 Assesses breathing for no more than 10 seconds. 	"Person is breathing normally."	
Obtains Consent	► Introduces self and asks, "May I help you?"	"Person consents."	
Assesses for Life- Threatening Conditions	 Quickly scans the person for life-threatening conditions. 	"The person is visibly shaking and pressing their hand over an obvious thigh wound. Blood is leaking around their hand and dripping onto the floor."	
Immediately Provides Appropriate First Aid	 Exposes the wound. Applies a single stack of 10 4" x 4" gauze pads over wound. Applies direct manual pressure over the wound using the heel of one hand with the other hand stacked on top of the first or the pads of 3 fingers of each of hand stacked on top of each other. 	"Blood is flowing contin- uously from the wound."	
Maintains and Increases Pressure	 Increases direct pressure over the wound. Doesn't remove pressure to add more gauze and doesn't remove blood-soaked materials. 	[Instructor, after about another 5 seconds] "The bleeding has stopped."	
Applies Bandage Over Dressing	 Wraps an elastic or self-adhesive roller bandage firmly over the gauze to help maintain pressure. 		

Procedure	Provider Action (Performance Criteria)	Instructor Prompt	Check Off
Removes PPE	 Correctly removes and disposes of gloves. Verbalizes washing hands immediately after removing gloves. 	[Instructor] "EMS arrives and takes over. Please demonstrate how to remove your gloves and what actions you would take next."	
	END PERFORMANCE EVALUATION		
☐ Successfully (Completed.	equired.	
Signature of HSI	Authorized Instructor HSI Instru	uctor Registry Number:	

Til.	
h	CI
8 1	101

performance evaluation five

ADULT FIRST AID - SEVERE ALLERGIC REACTION



Student Name	
	Class Date

You are a trained first aid provider responding to a call for help received over your workplace mobile device for a team member with trouble breathing. As you reach the scene, you see a person sitting at a break table. You have appropriate PPE and a first aid kit. Demonstrate what actions you would take next.

Procedure	Provider Action (Performance Criteria)	Instructor Prompt	Che
Performs Assessment	 Assesses scene safety. Takes (or verbalizes taking) standard precautions. Assesses responsiveness. Activates EMS and/or EAP. After activating, and unless they are readily available, sends someone to get the first aid kit and an AED. 	"The scene is safe." "Person is responsive." "EMS/EAP activated." "Other first aid providers are on the way, bringing an AED and an additional first aid kit."	
Assesses Breathing	 Assesses breathing for no more than 10 seconds. 	"Person is responsive and breathing."	
Obtains Consent	► Introduces self and asks, "May I help you?"	"Person consents."	
Assesses for Life- Threatening Conditions	 Quickly scans the person for life-threatening conditions. 	"The person is in obvious distress - trembling, wheezing, and coughing. You see swelling of their lips, eyelids, and face. With difficulty the person says, "Nut allergiesatepretzel bite had peanut butter inside."	
mmediately Provides Appropriate First Aid	 Allows the person to find the most comfortable position in which to breathe. 	"Person wants to sit up."	
erforms a econdary ssessment	► Looks for medical identification jewelry.	"Person is wearing a Medical Alert Bracelet on wrist that states, 'Nut Allergy - Use EpiPen®.'" [Instructor Note: Hand student epinephrine autoinjector trainer. Say "Here's their EpiPen®. They forgot how to use it."	

Procedure	Provider Action (Performance Criteria)	Instructor Prompt	Check Off
Correctly Uses Epinephrine Autoinjector Trainer	 Grasps epinephrine autoinjector trainer in fist with the orange tip pointing downward. Removes blue safety release. Positions trainer near middle of person's outer thigh. Swings and firmly pushes orange tip at a 90-degree angle against thigh until it clicks. Holds device firmly on thigh for 3 seconds. Removes the trainer from the thigh and massages the injection area for 10 seconds. 	[Instructor Note: Use instructions at the left are for the correct use of an EpiPen® epinephrine autoinjector trainer. If using a different manufacturer's training device, students should follow the manufacturer's instructions for use].	
	END PERFORMANCE EVALUATION		
☐ Successfully Co	mpleted. Not Successfully Completed. Remediation Re	equired.	
Signature of HSI A	uthorized Instructor HSI Instru	ictor Registry Number:	